



American Association for Cancer Research®

FINDING CURES TOGETHER®

**Embargoed for Release:** 8:30 a.m. ET, April 10, 2021

To interview Anna Plym, please contact Julia Gunther at [julia.gunther@aacr.org](mailto:julia.gunther@aacr.org) or 770-403-7690. For a photo of Plym, click [here](#). Visit our [newsroom](#).

## Healthy Lifestyle May Offset Risk of Lethal Prostate Cancer in Men With High Genetic Risk

PHILADELPHIA –Men who had a high genetic risk of developing prostate cancer were less likely to develop a lethal form of the disease if they maintained a healthy lifestyle, according to results presented during Week 1 of the virtual [AACR Annual Meeting 2021](#), held April 10-15.

[Prostate cancer](#) is the most common non-skin cancer detected in men, and the second leading cause of cancer deaths in men, after lung cancer. Genetic factors account for approximately 58 percent of the variability in prostate cancer risk, explained one of the study's lead authors, [Anna Plym, PhD](#), a postdoctoral research fellow at Brigham and Women's Hospital and Harvard T. H. Chan School of Public Health.

In this study, Plym and colleagues sought to evaluate whether increased genetic risk of prostate cancer could be offset by following a healthy lifestyle. Using a validated polygenic risk score for overall prostate cancer, the researchers quantified the genetic risk of prostate cancer in 10,443 men in the Health Professionals Follow-up Study for whom genotype data was available. They applied a validated lifestyle score for lethal prostate cancer which encompassed healthy weight, vigorous physical activity, not smoking, and high consumption of tomatoes, fatty fish, and reduced intake of processed meat.

In a median follow-up period of 18 years for overall prostate cancer and 22 years for lethal prostate cancer, the researchers identified 2,111 overall prostate cancer cases and 238 lethal prostate cancer cases. The study showed that men in the highest risk quartile according to their polygenic risk score were 5.4 times more likely to develop prostate cancer and 3.5 times more likely to develop lethal prostate cancer than those in the lowest risk quartile.

The researchers then measured the effect of adhering to a healthy lifestyle, and found that among men in the highest-risk quartile, those with the highest healthy lifestyle scores had about half the risk of developing lethal prostate cancer compared with those with the least healthy lifestyle. In this high-risk group, having a healthy lifestyle at the time of study entry was associated with a lifetime cumulative lethal prostate cancer incidence of 3 percent, compared with 6 percent for high-risk men having the least healthy lifestyle and 3 percent for the study population as a whole.

"The decreased risk of aggressive disease in those with a favorable lifestyle may suggest that the excess genetic risk of lethal prostate cancer could be offset by adhering to a healthy lifestyle," Plym said.

Adhering to a healthy lifestyle was not associated with a decreased risk of overall prostate cancer, nor did it affect men in the lower genetic risk quartiles. Plym said further research is necessary to determine why the benefit was limited to lethal prostate cancer risk in men with the highest genetic risk; one possible explanation is that the genetic variants that contribute to a high polygenic risk score are also the variants with the strongest interaction with lifestyle factors.

The study adds to a wide body of cancer prevention research that shows the benefit of a healthy lifestyle, including not smoking, maintaining a healthy weight, getting regular exercise, and eating a healthy diet.

Plym added that the results of this study underscore the importance of surveillance for those with a genetic predisposition to develop prostate cancer.

“Our findings add to current evidence suggesting that men with a high genetic risk may benefit from a targeted prostate cancer screening program, aiming at detecting a potentially lethal prostate cancer while it is still curable,” Plym said.

Plym noted that the study is observational, and therefore the association between healthy lifestyles and prostate cancer may not be a causal link.

This study was funded by the DiNovi Family Foundation, the National Cancer Institute at the National Institutes of Health, the William Casey Foundation, the Swedish Society for Medical Research, and the Prostate Cancer Foundation. Plym declares no conflicts of interest.

## **Abstract**

**Control Number:** 21-A-2868-AACR

**Session Type:** **Poster Session**

**Session Title:** **Gene-Gene and Gene-Environment Interactions**

**Session Time:** Saturday, April 10, 2021, 8:30 am - 11:59 pm

**Presentation Number:** 822

**Publishing Title:** Can the genetic risk of prostate cancer be attenuated by a healthy lifestyle?

**Author Block:** *Anna Plym, Yiwen Zhang, Konrad Stopsack, Bénédicte Delcoigne, Adam S. Kibel, Edward Giovannucci, Kathryn L. Penney, Lorelei A. Mucci.* Brigham and Women's Hospital, Boston, MA, Harvard T. H. Chan School of Public Health, Boston, MA, Memorial Sloan Kettering Cancer Center, New York, NY, Karolinska Institutet, Stockholm, Sweden

**Introduction:** Inherited genetic factors contribute significantly to prostate cancer risk and explains 58% of the variability in prostate cancer incidence. It is unclear if the increased genetic risk of prostate cancer, including progression to lethal disease, can be offset by adherence to a healthy lifestyle.

**Methods:** Using a validated polygenic risk score (PRS) for overall prostate cancer, we quantified the genetic risk of prostate cancer in 10,443 men in the Health Professionals Follow-up Study for whom genotype data was available. We applied a validated lifestyle score for lethal prostate cancer (including healthy weight, vigorous physical activity, not smoking, and high consumption of tomatoes, fatty fish, and reduced intake of processed meat) and examined the incidence of overall and lethal (metastatic disease or prostate cancer-specific death) prostate cancer from the date of blood (1993-1999) or cheek collection (2005-2006) through 2014 (2016 for lethal disease).

Multivariable Cox proportional hazards models were used to estimate the risk of overall and lethal prostate cancer by joint categories of genetic risk (PRS quartiles) and a time-varying lifestyle score (1-2: least healthy, 3: moderate healthy, and 4-6: most healthy). Both unweighted and inverse probability weighted (IPW) models (to account for possible bias arising from the genotype sampling design) were applied. Lifetime cumulative incidence was estimated using regression standardization.

**Results:** We observed 2,111 prostate cancer and 238 lethal prostate cancer events during a median follow-up of 18 and 22 years, respectively. The PRS enabled risk stratification for both overall and lethal prostate cancer, with men in the highest genetic risk quartile having a 5.4-fold increased risk of overall prostate cancer ( $HR_{ipw} = 5.39$ , 95% CI = 4.59-6.33) and a 3.5-fold increased risk of lethal prostate cancer ( $HR_{ipw} = 3.53$ , 95% CI = 2.34-5.32) compared with men in the lowest genetic risk quartile. Among men in the highest genetic risk quartile, adhering to a healthy lifestyle was significantly associated with a decreased risk of lethal prostate cancer ( $HR_{ipw} = 0.54$ , 95% CI = 0.31-0.94) compared with the least healthy lifestyle. Adhering to healthy lifestyle was not associated with a decreased risk of overall prostate cancer ( $HR_{ipw} = 1.01$ , 95% CI = 0.84-1.22). In the group of men with highest genetic risk, having a healthy lifestyle at study entry was associated with a lifetime

## **Abstract**

### **Body:**

cumulative incidence of lethal prostate cancer of 3%, lower than for men having the least healthy lifestyle (6%) and similar to the population average (3%).

**Conclusion:** In this large prospective cohort of US men, inherited genetic factors increased the risk of both overall and lethal prostate cancer. The associated decreased risk of aggressive disease in those with a favorable lifestyle may suggest that the excess genetic risk of lethal prostate cancer could be offset by adhering to a healthy lifestyle.

###

Follow the meeting on Twitter at [#AACR21](#)

Follow us: [Cancer Research Catalyst](#); [Twitter](#); and [Facebook](#)

For AACR information, visit [Fast Facts](#).

#### **About the American Association for Cancer Research**

Founded in 1907, the American Association for Cancer Research (AACR) is the world's first and largest professional organization dedicated to advancing cancer research and its mission to prevent and cure cancer. AACR membership includes 48,000 laboratory, translational, and clinical researchers; population scientists; other health care professionals; and patient advocates residing in 127 countries. The AACR marshals the full spectrum of expertise of the cancer community to accelerate progress in the prevention, diagnosis, and treatment of cancer by annually convening more than 30 conferences and educational workshops—the largest of which is the AACR Annual Meeting, with more than 74,000 attendees for the 2020 virtual meetings and more than 22,500 attendees for past in-person meetings. In addition, the AACR publishes nine prestigious, peer-reviewed scientific journals and a magazine for cancer survivors, patients, and their caregivers. The AACR funds meritorious research directly as well as in cooperation with numerous cancer organizations. As the Scientific Partner of Stand Up To Cancer, the AACR provides expert peer review, grants administration, and scientific oversight of team science and individual investigator grants in cancer research that have the potential for near-term patient benefit. The AACR actively communicates with legislators and other policymakers about the value of cancer research and related biomedical science in saving lives from cancer. For more information about the AACR, visit [www.AACR.org](http://www.AACR.org).